

Product Description

BspE I is a Type IIP restriction endonuclease expressed in *Escherichia coli* carrying the BspE I gene cloned from *Bacillus*.

Its recognition and cleavage sequences are shown below:

5'...T ↓ CCGGA...3'

3'...AGGCC ↑ T...5'

Components

Components	BR1G106-01 1,000 U	BR1G106-02 2,000 U	BR1G106-03 5,000 U
BspE I (10 U/μL)	1×0.1 mL	1×0.2 mL	1×0.5 mL
10× Cut-Buffer	1×1 mL	1×1 mL	3×1 mL

Unit Definition

One unit is defined as the amount of enzyme required to completely and specifically cleave 1 μg of internal control plasmid template in 1 hour at 37°C.

Storage

Store at -20±5°C.

Notes

1. For research use only. Not for use in clinical diagnosis.
2. Mix thoroughly before use. Avoid repeated freeze-thaw cycles.
3. The volume of restriction endonuclease added should not exceed one-tenth of the total reaction volume.
4. Suitable for digestion of genomic DNA, plasmid DNA, and PCR products.
5. Digestion for longer than 3 h is not recommended, as prolonged incubation may lead to star activity.

Prepare Reaction Mix

Prepare the reaction mix on ice as follows:

Components	Volume per Reaction	Concentration in Master Mix
10× Cut-Buffer	5 μL	1×
BspE I (10 U/μL)	1 μL	0.2 U/μL
DNA Substrate	1 μg	—
ddH ₂ O	To 50 μL	—

Reaction Programe

Incubate at 37°C for 1 h. Heat inactivation: incubate at 80°C for 20 min.